

Linear Enamel Hypoplasia and Age-at-Death at Medieval (11th–16th Centuries) St. Gregory's Priory and Cemetery, Canterbury, UK

JUSTYNA JOLANTA MISZKIEWICZ*

School of Anthropology and Conservation, University of Kent, Marlowe Building, Canterbury, CT2 7NR, United Kingdom

ABSTRACT

Linear enamel hypoplasia (LEH) is a macroscopically detectable band-like dental defect, which represents localized decrease in enamel thickness caused by some form of disruption to a child's health. Such dental deformations are utilized in osteoarchaeological research as permanent markers of childhood physiological stress and have been extensively studied in numerous ancient human populations. However, currently there is no such data for medieval populations from Canterbury, UK. Here, LEH is examined in the context of age-at-death in human burials from the medieval St. Gregory's Priory and adjacent cemetery (11th–16th centuries), Canterbury, UK. The cemetery and Priory burials represented lower (n = 30) and higher status (n = 19) social groups, respectively.

Linear enamel hypoplastic defects were counted on mandibular and maxillary anterior permanent teeth (n = 374). The age and sex of each skeleton were estimated using standard methods. Differences in LEH counts, age-at-death, and LEH formation ages were sought between the two social groups. Results indicate significantly greater frequencies of LEH in the Cemetery (mean = 17.6) compared to the Priory (mean = 7.9; t = 3.03, df = 46, p = 0.002). Adult age-at-death was also significantly lower in the Cemetery (mean = 39.8, df = 47, p = 0.013). Hypoplasia formation years) compared to the Priory burials (mean = 44.1 years; ages differed significantly between the Priory (mean = 2.49 years) and Cemetery (mean = 3.22 years; t = 2.076; df = 47; p = 0.034) individuals.

Results indicate that childhood stress may reflect adult mortality in this sample, and that the wellbeing of individuals from diverse social backgrounds can be successfully assessed using LEH analyses. Results are discussed in terms of the multifactorial aetiology of LEH, as well as weaning-related LEH formation.

Key words: linear enamel hypoplasia; age-at-death; social status; medieval Canterbury

Introduction

Childhood physiological stress can disturb enamel formation, producing localized hypoplastic defects that can emerge on the outer tooth surface (Hillson, 1992; Larsen, 1997; Reid & Dean, 2000, 2006; Hubbard et al., 2009). These malformations occur in several forms (e.g. irregular depressions, pitting, or furrowing) (Hillson & Bond, 1997), but linear enamel hypoplasia (LEH) is the most commonly reported (Hillson, 1996; Griffin & Donlon, 2008). The health of archaeological samples of modern human populations can be deduced from LEH analyses (Larsen, 1997; Roberts & Manchester, 2005), because of the non-specific nature of this dental pathology (Goodman & Armelagos, 1985). Although many aetiological explanations, such as childhood disease (e.g. measles) and malnutrition (Goodman & Armelagos, 1988; Larsen, 1997; Roberts & Manchester, 2005), as well as weaning (Blakey et al., 1994), and accidental consumption of toxins (Neiburger, 1990) have been put forward as factors underlying LEH formation, it can be difficult to identify these causes in human archaeological samples. Thus, the majority of prior research has focused on LEH as a permanent marker of 'general' childhood physiological stress (e.g. Rose et al., 1978; Goodman & Armelagos, 1988; Wright, 1997; Palubeckaitė, 2001; Šlaus et al., 2002; King et al., 2005; Boldsen, 2007; Griffin & Donlon, 2007; Mays & Barber, 2008; Dabbs, 2011; Krenz-Niedbała & Kozłowski, 2011).

Numerous studies have examined the relationship between hypoplastic defects and stress in the context of social status from archaeological sites (e.g. Lanphear, 1990; Cucina and Işcan, 1997; Cucina, 2002; Starling & Stock, 2007; Tomczyk et al., 2007; Mayes & Barber, 2008; Dabbs, 2011). A variety of themes have been covered, including: LEH prevalence correlated with weaning and differences in subsistence strategies (Lanphear, 1990); varied population stress levels associated with cultural amelioration (Dabbs, 2011); high status or more 'privileged' ancient groups affected by LEH (Cucina and Işcan, 1997; Mayes & Barber, 2008); or links between LEH and periods of food scarcity (Tomczyk et al., 2007). When examined in the context of age-at-death, LEH records also show positive associations, implying that severe childhood stress can affect longevity (e.g. Šlaus et al., 2002; Palubeckaitė et al., 2002; King et al., 2005; Boldsen, 2007; Armelagos et al., 2009). However, not all projects implement consistent methodological controls when collecting LEH data. For example, in order to avoid bias and increase sample size some studies choose to use only one tooth (often from one side of the jaw) (e.g. Boldsen, 2007), but this restricts addressing potential systemic stress. Often, more specific tooth types (e.g. anterior dentition only) or jaw sides are chosen and control for dental calculus, but not wear is implemented (or at least reported) (e.g. Šlaus et al., 2002; Palubeckaitė et al., 2002). Only a few studies report complete exclusion of heavily worn dentition and analyze specific tooth types that display minimal attrition and dental calculus (in some cases, the latter is manually removed for the purpose of analysis) (e.g. Cucina, 2002; Griffin & Donlon, 2007). When examining stress markers registered in teeth, it is indeed of importance to assign as many methodological controls as possible to account for age-related tooth loss and dental wear, and also identify systemic stress by comparing more than one tooth type (e.g. King et al., 2005), particularly when two or more groups are contrasted.

It is clear that osteoarchaeology has seen an abundance of research analyzing LEH in relation to other factors in a variety of past populations. However, no prior published analyses considered addressing (potential) childhood stress and possibly linked social status in human groups of medieval Canterbury (Kent, UK) (Figure 1). Historical aspects of medieval British societies allow constructing hypotheses that aim to study any links between health and stress as the Middle Ages were dominated by feudalism and the church, causing populations to divide into different socio-economic 'classes' (Hull, 1989). Consequently, these were reflected in the quality of life led by individuals representing different status groups, as demonstrated in the accelerated spread of infectious disease (such as leprosy) (Roberts & Manchester, 2005) and restricted access to exclusive types of foods (Dyer, 2003) in lower ranking individuals (Hull, 1989). Thus, results of a carefully controlled study carried out on two distinct archaeological groups, dated to the medieval period (11th – 16th centuries) from Canterbury (UK) are here reported. It is hypothesized

that adult age-at-death estimates, LEH counts, and LEH age formation estimates can aid the identification of stress and mortality caused by differences in social status.

Materials and methods

Study population

Medieval human skeletal remains recovered at the St. Gregory's Priory and a nearby Cemetery in the Northgate area of Canterbury (Kent, UK) (Figure 1), were examined. St. Gregory's Priory was established in the 11th century by Lanfranc, the Archbishop of Canterbury, and it functioned until the early 16th century (Woodcock, 1956:xi; Tatton-Brown, 1995; Hicks & Hicks, 2001). Historical and archaeological textual records imply that there was a distinct divide in the socio-economic status between individuals buried within the cemetery and those occupying St. Gregory's Priory (Tatton-Brown, 1995; Hicks & Hicks, 2001). The Priory was established for the clergy to minister to the sick and provide burial for financially depleted members of the medieval society in the associated cemetery. Therefore, burials examined in this study represented two distinct social groups, higher (Priory n=19) and lower status (Cemetery n=30), as inferred from burial location (Hicks & Hicks, 2001). The Priory originally contained a considerably small amount of burials due to its historical context. Furthermore, excluding juvenile remains and applying restrictive dental controls means that this study evaluates a limited sample size, but on the other hand, it presents a carefully controlled methodological approach. Osteological and statistical procedures

Adult age-at-death and sex were estimated following standard osteological methods (Buikstra & Ubelaker, 1994). In total, 374 (maxillary and mandibular) permanent canines and incisors were examined, as approximately 4.6% teeth were missing due to ante- or post-mortem loss as identified from empty tooth sockets. Posterior tooth rows were excluded from analyses, because they exhibit age-related occlusal attrition in adult samples (Miles, 1962; Hillson, 1996). Focusing on anterior dentition allowed for comparisons of the same tooth types across the board. In order to ensure that anterior wear did not skew results by erasing evidence of any formed defects, only comparably worn teeth displaying minimal amount of calculus (i.e. unmasking hypoplastic defects) were examined, further reducing sample size. This was achieved by ensuring that only maximum one third of the occlusal tooth section (after intermediate, preceded by cervical) was worn down in all studied individuals.

Hypoplastic defects were counted following the 'Field method' (Hassett, 2011) (Figure 2), which involves identifying lines macroscopically (Buikstra & Ubelaker, 1994; Brickley & McKinley, 2004). The anterior surface of each tooth with detected LEH was examined using a standard magnifying glass to minimize under- or over-estimating of LEH total counts for each tooth. A hypoplastic line was counted only if it extended horizontally across the labial tooth surface, measuring a minimum three quarters in length (e.g. Griffin & Donlon, 2008). Following this, the location of lines was compared to the timings of anterior tooth growth diagrams devised by Reid and Dean (2000; 2006) to estimate the age of LEH formation. Although many prior LEH publications have used other sources of LEH age determination, such as tooth mineralization stages by Bass (1995) (e.g. Boldsen, 2007) or a more conventional method by Massler et al. (1941) (e.g. Palubeckaitė et al., 2002; Krenz-Niedbala & Kozłowski, 2011); Reid and Dean's (2000; 2006), work on tooth formation times is based on histological examinations that take into account the so-called hidden (appositional) enamel (King et al., 2002) providing a reliable source for obtaining LEH formation age. A criterion identifying systemic stress was set to be confirmative if two or more teeth exhibited simultaneously forming LEH bands (e.g. Palubeckaitė et al., 2002; King et al., 2005).

Statistical analyses were performed in SPSS/PASW 18.0. It was impossible to explore sex differences as female skeletons were a minority in each group. This has resulted in pooling sexes together. Due to high and comparable records of presence of LEH in both groups, no statistical tests were run to determine disparities in prevalence as identified by dental defect presence or absence, rather differences in the frequencies (e.g. Boldsen, 2007; Griffin & Donlon, 2008) of hypoplasia and age-at-death, as well as hypoplastic line formation ages were sought using an independent samples t-test.

Results

Descriptive statistics are available in Tables 1–3. Presence of hypoplasia was recorded in the vast majority of individuals, with only two samples from the cemetery and four from the Priory displaying entirely unaffected dentition. The highest counts of lines were noted for upper left central incisor and both lower canines in the Cemetery (Table 2), whereas individuals from the Priory exhibited the highest numbers of hypoplastic lines on the upper left central incisor and lower right canine (Table 2). More hypoplastic bands were recorded among the Cemetery individuals (mean =17.6) than those from the Priory (mean =7.9) (Table 3). An independent samples t-test produced statistically significant results ($t = 3.03$, $df = 45.815$, $p = .002$, one-tailed). Age-at-death estimates differed significantly between Cemetery (mean =39.8 years) and Priory (mean =44.1 years) samples ($t = 2.275$, $df = 47$, $p = 0.0135$, one-tailed) (Table 1). Finally,

Discussion

The results from this study support the notion that a low socio-economic status can negatively affect an individual's wellbeing and lead to a disrupted enamel formation during childhood, and consequently shortened longevity. All hypoplastic bands recorded in this study reflected regular patterns, with multiple simultaneously developing lines frequently displayed in both groups. This regularity is suggestive of prevailing physiological alterations (Goodman et al., 1988; Hillson, 1992; 1996), but the data also indicates experiences of systemic stress (King et al., 2005). One individual line could be inferred to have been associated with a single traumatic event, having no relationship with social status or continual stress (Malville, 1997).

Both groups were almost completely affected by LEH; however, the Cemetery individuals displayed significantly higher LEH frequencies, implying that their health was more heavily disrupted during childhood years. It is unlikely that individuals from the Priory did not produce hypoplastic lines, because forming enamel is highly sensitive to many forms of stressors (May et al., 1993; Goodman & Rose, 1990). In this study, Priory individuals can be inferred to have been exposed to a lesser extent of stress when compared to the Cemetery group. Thus, confirming that LEH affects the dentition of more privileged, as well as lower ranking human groups, in line with LEH presence outcomes reported in studies analyzing high status archaeological human samples (e.g. Cucina & Işcan, 1997; Palubeckaitė et al., 2002; Mayes & Barber, 2008).

The Priory individuals had a mean hypoplasia formation age of 2.49 years – a much earlier result than in the Cemetery (3.22 years). It could be tempting to propose that dietary implications of weaning and post-weaning periods which often lead to nutritional imbalances in the early years of childhood (Goodman & Rose, 1990; Blakey et al., 1994; Kennedy, 2005) could have contributed to the formation of LEH in this group, because (on average) human juveniles undergo this event around 2.5 years of age (Kennedy, 2005). It is stressed that in this article, 'weaning' is defined as the cessation of breast-feeding and inclusion of solid non-maternal food sources into juvenile diet (World Health Organization Website, 2002; Griffiths et al., 2007). Although the age at which non-maternal food sources are introduced to weaned children is highly variable amongst human societies (e.g. Stewart-Macadam & Dettwyler, 1995; Griffiths et al., 2007), many osteoarchaeologists assign weaning related significance to 'peak' LEH records (e.g. Lanphear, 1990; Iregren, 1992; Ubelaker, 1992; Moggi-Cecchi et al., 1994; Wright, 1997). On the other hand, some studies have shown that such a link is weak or possibly non-existent (e.g. Blakey et al., 1994; Corruccini et al., 1985; Santos & Coimbra, 1999; Saunders & Keenleyside, 1999; Wood, 1996). Thus, the most suitable interpretation of the above result would be achieved if data informative of the weaning age for this population from Canterbury was available. Unfortunately, no such background information can be found in the literature. Therefore, the reported differences in LEH formation ages remain attributed to non-specific physiological disruptions specific to social status.

To further support the link between health and social status, this project predicted that age-at-death estimates would differ significantly between the medieval Cemetery and Priory. The results imply that longevity records in this sample were in line with the documented differences in status. Mean age-at-death

estimates for the former and the latter were 39.8 and 44.1 years, respectively, still falling within the ‘middle adult’ age category (35–50 years) (Buikstra & Ubelaker, 1994), but were evidently significantly different between each other. Perhaps, more pronounced results would have been obtained had sub-adult and juvenile burials been considered in the analysis. It is potentially a future step for studies exploring the health of this medieval population. Nevertheless, the results do reflect outcomes from other studies that addressed the relationship between childhood stress and longevity (e.g. Šlaus et al., 2002; Palubeckaitė et al., 2002; Steckel, 2005; Boldsen, 2007). Additionally, the present study further adds to Armelagos and colleagues’ (2009) bioarchaeological support for the Barker hypothesis, which proposes that pathological events occurring early in life make a negative impact on the adult health (Paneth et al., 1996).

The aetiology of LEH and definition of stress in a bioarchaeological context are of a multi-faceted nature (Goodman & Armelagos, 1988; Neiburger, 1990; Ribot & Roberts, 1996). Researchers have argued that hypoplastic lines are not necessarily valid indicators of physiological (particularly malnutrition) stress (e.g. Neiburger, 1990). Several types of causation of enamel defects have been proposed, including genetic predisposition for defective enamel; single traumatic events; poisoning with chemical or other harmful substances; even cultural modification of teeth (Neiburger, 1990). These are all valid points; however, the degree to which such aspects would influence population-based health profiles is relatively low, as high frequency and severity of LEH would still reflect a continuum of stressful events in a child’s life.

Finally, this study finds that hypoplastic defects are valid indicators of stress, especially in the context of age-at-death, when analyzed from a social status hierarchy perspective. However, a purely macroscopic approach to LEH examination could be under-estimating pathological structure of enamel in affected dentition (Hassett, 2011); hence, careful inferences are made about this ancient population. Here, differences in the examined variables indicate exposure to uncertain conditions and suggest continuous physiological disruptions faced during childhood in the Cemetery individuals. Teeth remain the most useful ancient skeletal specimens for detecting markers of stress, as other bone pathologies acquired during childhood are generally difficult to diagnose in adult samples, due to bone modelling and remodelling and thus the disappearance of lesions during bone growth (McHenry & Schulz, 1976; Roberts & Manchester, 2005). The enamel’s ability to register disruptive events was useful in researching childhood stress in this sample, particularly as it reflected textually documented social status divide of this population.

Conclusions

The results indicate that childhood stress may reflect adult mortality in this sample, and that the wellbeing of individuals from diverse social backgrounds can be successfully examined via carefully controlled LEH studies. Frequent stress can be assumed for Cemetery individuals, as inferred from the presence of multiple and regular LEH lines. Both groups display LEH; however, significantly lower frequencies in the Priors sample suggest stress of a lesser extent. The mean age-at-death estimates fell within the ‘middle adult’ category (35–50 years) (Buikstra & Ubelaker, 1994). Perhaps, biological significance of social status would be more pronounced if juvenile burial frequencies were considered. This study provides quantitative data that supports the social status divide of this British medieval population.

Acknowledgments

The author is grateful to the editor and three reviewers whose invaluable comments and suggestions helped improve this article. Thanks go to Patrick Mahoney for his supervision throughout this project, Chris Deter for her remarks on the earlier version of this manuscript, and the Biological Anthropology Research Laboratory at the University of Kent for allowing access to samples.

References

- Armstrong GJ, Goodman AH, Harper KN, Blakey ML. 2009. Enamel hypoplasia and early mortality: Bioarchaeological support for the Barker hypothesis. *Evolutionary Anthropology: Issues, News, and Reviews*, 18: 261–271. DOI: 10.1002/evan.20239
- Bass WM. 1995. *Human Osteology: A Laboratory and Field Manual*. Columbia, MO: Missouri Archaeological Society. Special Publication No. 2.
- Blakey ML, Leslie TE, Reidy JP. 1994. Frequency and chronological distribution of dental enamel hypoplasia in enslaved Americans: a test of the weaning hypothesis. *American Journal of Physical Anthropology* 95: 371–383. DOI: 10.1002/ajpa.1330950402
- Boldsen JL. 2007. Early childhood stress and adult age mortality – a study of dental enamel hypoplasia in the Medieval Danish village of Tirup. *American Journal of Physical Anthropology* 132: 59–66. DOI: 10.1002/ajpa.20467
- Brickley ML, McKinley JI. 2004. Guidelines to standards for recording human remains. British Association for Biological Anthropology and Osteoarchaeology and Institute of Field Archaeologists: Southampton and Whitenights.
- Buikstra JE, Ubelaker DH. 1994. Standards for Data Collection from Human Skeletal Remains. Fayetteville, Arkansas: Arkansas Archaeological Survey Report 44.
- Cucina A, İşcan MY. 1997. Assessment of enamel hypoplasia in a high status burial site. *American Journal of Human Biology* 9: 213–222. DOI: 10.1002/(SICI)1520-6300(1997)9:2<213:: AID-AJHB7>3.0.CO;2-0
- Cucina A. 2002. Brief communication: Diachronic investigation of linear enamel hypoplasia in prehistoric skeletal samples from Trentino, Italy. *American Journal of Physical Anthropology* 119: 283–287. DOI: 10.1002/ajpa.10135.
- Corruccini RS, Handler JS, Jacobi KP. 1985. Chronological distribution of enamel hypoplasias and weaning in a Caribbean slave population. *Human Biology* 57: 699–711.
- Dabbs GR. 2011. Health status among prehistoric Eskimos from Point Hope, Alaska. *American Journal of Physical Anthropology*, 146: 94–103. DOI: 10.1002/ajpa.21556
- Dyer C. 2003. Peasants, c. 1100- c.1315. In *Making a Living in the Middle Ages: The People of Britain 850–1520*, C Dyer (ed.). Penguin Books: London; 155–186.
- Goodman AH, Armstrong GJ. 1985. Factors affecting the distribution of enamel hypoplasias within the human permanent dentition. *American Journal of Physical Anthropology* 68: 479–493. DOI: 10.1002/ajpa.1330680404
- Goodman AH, Armstrong GJ. 1988. Childhood stress and decreased longevity in a prehistoric population. *American Anthropologist* 90: 936–944. DOI: 10.1525/aa.1988.90.4.02a00120
- Goodman AH, Rose JC. 1990. Assessment of systemic physiological perturbations from dental enamel hypoplasias and associated histological structures. *Yearbook of Physiological Anthropology* 33: 59–110. DOI: 10.1002/ajpa.1330330506
- Griffin RC, Donlon D. 2007. Dental enamel hypoplasias and health changes in the Middle Bronze Age – Early Iron Age transition at Pella in Jordan. *Journal of Comparative Human Biology* 58: 211–220. DOI:10.1016/j.jchb.2006.08.006
- Griffin RC, Donlon D. 2008. Patterns in dental enamel hypoplasia by sex and age at death in two archaeological populations. *Archives of Oral Biology* 54: S93-S100. DOI:10.1016/j.archoralbio.2008.09.012

- Griffiths LJ, Tate AR, Dezateux C, Millenium Cohort Study Child Health Group. 2007. Do early infant feeding practices vary by maternal ethnic group? *Public Health Nutrition* 10: 957–964. DOI: 10.1017/S1368980007665513
- Hassett BR. 2011. Evaluating sources of variation in the identification of linear hypoplastic defects of enamel: a new quantified method. *Journal of Archaeological Science*, 39: 560–565. DOI: 10.1016/j.jas.2011.10.017
- Hicks M, Hicks A. 2001. *St. Gregory's Priory, Northgate, Canterbury Excavations 1988–1991*. Canterbury Archaeological Trust Ltd: Volume II.
- Hillson SW. 1992. Dental enamel growth, perikymata and hypoplasia in ancient tooth crowns. *Journal of the Royal Society of Medicine* 85: 460–466.
- Hillson S. 1996. *Dental Anthropology*. Cambridge University Press: Cambridge.
- Hillson S, Bond S. 1997. Relationship of enamel hypoplasia to the pattern of tooth crown growth: a discussion. *American Journal of Physical Anthropologists* 104: 89–103. DOI:10.1002/(SICI)1096-8644(199709)104:1<89::AID-AJPA63.0.CO;2-8
- Hubbard A, Guatelli-Steinberg D, Sciulli PW. 2009. Under restrictive conditions, can the widths of linear enamel hypoplasias be used as relative indicators of stress episode duration? *American Journal of Physical Anthropology* 138: 177–189. DOI: 10.1002/ajpa.20917
- Hull BB. 1989. Religion, afterlife, and property rights in the high Middle Ages. *Studies in Economic Analysis* 12: 3–21.
- Iregren E. 1992. Scandinavian women during the medieval period; health, childbirth and childcare. *Collegium Anthropologicum* 16: 59–82.
- Kennedy GE. 2005. From the ape's dilemma to the weanling's dilemma: early weaning and its evolutionary context. *Journal of Human Evolution* 48: 123–145. DOI: 10.1016/j.jhevol.2004.09.005
- King T, Humphrey LT, Hillson S. 2005. Linear enamel hypoplasias as indicators of systemic physiological stress: evidence from two known age-at-death and sex populations from postmedieval London. *American Journal of Physical Anthropology* 128: 547–559. DOI: 10.1002/ajpa.20232
- King T, Hillson S, Humphrey LT. 2002. A detailed study of enamel hypoplasia in a post-medieval adolescent of known age and sex. *Archives of Oral Biology* 47: 29–39. DOI:10.1016/S0003-9969(01)00091-7
- Krenz-Niedbała M, Kozłowski T. 2011. Comparing the chronological distribution of enamel hypoplasia in Rogowo, Poland (2nd century AD) using two methods of defect timing estimation. *International Journal of Osteoarchaeology* DOI: 10.1002/oa.1262
- Larsen CS. 1997. Stress and deprivation during the years of growth and development and adulthood. In *Bioarchaeology: Interpreting Behaviour from the Human Skeleton*, CS Larsen (ed.). Cambridge University Press: United Kingdom; 6–63.
- Lanphear KM. 1990. Frequency and distribution of enamel hypoplasias in a historic skeletal sample. *American Journal of Physical Anthropology* 81: 35–43. DOI: 10.1002/ajpa.1330810106
- May RL, Goodman AH, Meindl RS. 1993. Response of bone and enamel formation to nutritional supplementation and morbidity among malnourished Guatemalan children. *American Journal of Physical Anthropology* 92: 37–51. DOI: 10.1002/ajpa.1330920104
- Malville NJ. 1997. Enamel hypoplasia in ancestral Puebloan populations from southwestern Colorado. 1. Permanent dentition. *American Journal of Physical Anthropology* 102: 351–367. DOI: 10.1002/(SICI)1096-8644(199703)102:3<351:AID-AJPA5>3.0.CO;2-Y
- Massler M, Schour I, Poncher HG. 1941. Developmental pattern of the child as reflected in the calcification pattern of the teeth. *American Journal of Disease of Children* 62: 33–67.

- Mayes AT, Barber SB. 2008. Osteobiography of a highstatus burial from the lower Rio Verde Valley of Oaxaca, Mexico. *International Journal of Osteoarchaeology* 18: 573–588. DOI: 10.1002/oa.1011
- Miles AEW. 1962. Assessment of the ages of a population of Anglo-Saxons from their dentitions. *Proceedings of the Royal Society of Medicine* 55: 881–886.
- McHenry HM, Schulz PD. 1976. The association between Harris Lines and enamel hypoplasia in prehistoric California Indians. *American Journal of Physical Anthropology* 44: 507–512. DOI: 10.1002/ajpa.1330440313
- Moggi-Cecchi J, Pacciani E, Pinto-Cisternas J. 1994. Enamel hypoplasia and age at weaning in 19th century Florence, Italy. *American Journal of Physical Anthropology* 93: 299–306. Neiburger EJ. 1990. Enamel hypoplasias: Poor indicators of dietary stress. *American Journal of Physical Anthropology* 82: 231–233. DOI: 10.1002/ajpa.1330820211
- Palubeckaitė Z. 2001. Patterns of linear enamel hypoplasia in Lithuanian Iron Age population. *Variability and Evolution* 9: 75–87.
- Palubeckaitė Z, Jankauskas R, Boldsen J. 2002. Enamel hypoplasia in Danish and Lithuanian late Medieval/ early modern samples: a possible reflection of child morbidity and mortality patterns. *International Journal of Osteoarchaeology* 12: 189–201. DOI: 10.1002/oa.607
- Paneth N, Ahmed F, Stein A. 1996. Early nutritional origins of hypertension: a hypothesis still lacking support. *Journal of Hypertension* 14: S121–S129.
- Reid DJ, Dean MC. 2000. Brief Communication: The timing of linear hypoplasias on human anterior teeth. *International Journal of Physical Anthropology* 113: 135–139. DOI: 10.1002/1096-8644(200009)113:1 <135:AID-AJPA13 3.0.CO;2-A
- Reid DJ, Dean MC. 2006. Variation in modern human enamel formation times. *Journal of Human Evolution* 50: 329–346. DOI:10.1016/j.jhevol.2005.09.003
- Ribot I, Roberts CA. 1996. A study of non-specific stress indicators and skeletal growth in two medieval subadult populations. *Journal of Archaeological Science* 23: 67–79.
- Roberts CA, Manchester K. 2005. *The Archaeology of Disease*. Sutton Publishing: Stroud. ISBN 0 7509 2683 X.
- Rose JC, Armelagos GJ, Lallo JW. 1978. Histological enamel indicator of childhood stress in prehistoric skeletal samples. *American Journal of Physical Anthropology* 49: 511–516. DOI: 10.1002/ajpa.1330490411.
- Santos RV, Coimbra CE, Jr. 1999. Hardships of contact: enamel hypoplasias in Tupi-Monde Amerindians from the Brazilian Amazonia. *American Journal of Physical Anthropology* 109: 111–127.
- Saunders S, Keenleyside A. 1999. Enamel hypoplasia in a Canadian historic sample. *American Journal of Human Biology* 11: 513–524.
- Šlaus M, Kollmann D, Novak SA, Novak M. 2002. Temporal trends in demographic profiles and stress levels in Medieval (6th – 13th Century) population samples from continental Croatia. *Basic Sciences* 43: 598–605.
- Starling AP, Stock JT. 2007. Dental indicators of health and stress in early Egyptian and Nubian agriculturalists: a difficult transition and gradual recovery. *American Journal of Physical Anthropology* 134: 520–528. DOI: 10.1002/ajpa.20700
- Steckel RH. 2005. Young adult mortality following severe physiological stress in childhood: skeletal evidence. *Economics and Human Biology* 3: 314–328. DOI:10.1016/j.ehb.2005.05.006
- Stewart-Macadam P, Dettwyler KA. 1995. *Breastfeeding: Biocultural perspectives*. Walter de Gruyter: New York.

- Tatton-Brown T. 1995. The beginnings of St. Gregory's Priory and St. John's Hospital in Canterbury. In *Canterbury and the Norman Conquest: Churches, Saints and Scholars, 1066–1109*, R Eales, R Sharpe (eds.). Hambledon Continuum: UK.
- Tomczyk J, Soltysiak A, Tomczyk-Gruca M. 2007. Temporal changes in frequency of enamel hypoplasia in the middle Euphrates Valley (Syria). *Human Biology Budapest* 30: 87–97.
- Ubelaker DH. 1992. Enamel hypoplasia in ancient Ecuador. In: *Recent Contributions to the Study of Enamel Developmental Defects*. *Journal of Paleopathology, Monographic Publication, 2*, AH Goodman, LL Capasso (eds.). Edigrafital: Teramo; 207–217.
- World Health Organization (WHO) 2002. *Infant and Young Child Nutrition. Global Strategy on Infant and Young Child Feeding*: Geneva.
- Woodcock AM. 1956. Introduction. In *Cartulary of the Priory of St. Gregory, Canterbury*. Offices of The Royal Historical Society: London; vii-xix.
- Wright LE. 1997. Intertooth patterns of hypoplasia expression: implications for childhood health in the classic Maya collapse. *American Journal of Physical Anthropology* 102: 233–247. DOI: 10.1002/(SICI)1096-8644(199702)102:2<233::AID-AJPA6>3.0.CO;2-Z
- Wood L. 1996. Frequency and chronological distribution of linear enamel hypoplasia in a North American colonial skeletal sample. *American Journal of Physical Anthropology* 100: 247–259.